

Induction of vascular endothelial growth factor release by transcutaneous frequency modulated neural stimulation in diabetic polyneuropathy

M. Bevilacqua¹, L.J. Dominguez², M. Barrella¹, and M. Barbagallo²

¹Endocrinology and Diabetes Unit and LORENZ Research Center, Department of Medicine, Luigi Sacco Hospital (Vialba)-University of Milan, Milan; ²Geriatric Unit, Department of Internal Medicine and

Reprinted from:

JOURNAL OF ENDOCRINOLOGICAL INVESTIGATION

JOURNAL OF ENDOCRINOLOGICAL INVESTIGATION

Official Journal of
Italian Society of Endocrinology
Italian Society of Medical Andrology
Italian Society of Pediatric Endocrinology and Diabetology



Editrice Kurtis s.r.l.
Via Luigi Zoja, 30
20153 Milano, Italy

<http://www.kurtis.it> - e-mail: jei@kurtis.it - Tel. +39 02 48202740

JOURNAL OF ENDOCRINOLOGICAL INVESTIGATION

Official Journal of
Italian Society of Endocrinology
Italian Society of Medical Andrology
Italian Society of Pediatric Endocrinology and Diabetology

Journal recognized as having "high cultural value"
by the Ministry of Cultural Heritage

General Information

The **Journal of Endocrinological Investigation** publishes original studies on clinical and experimental research in endocrinology and related fields. Details on the types of manuscripts that will be considered for publication are given in the Instructions to Authors. Full version of the Instructions to Authors is published in the January issue and can be downloaded from www.kurtis.it

All editorial correspondence should be addressed to the Editorial Office:

Editor-in-Chief
Ezio Ghigo MD
Division of Endocrinology
University of Turin
C.so Dogliotti, 14
10126 Turin, Italy
Tel. +39 011.6334317
Fax +39 011.6647421

E-mail: elisabetta.bersezio@unito.it

Editorial Office
Elisabetta Bersezio
Division of Endocrinology
University of Turin
C.so Dogliotti, 14
10126 Turin, Italy
Tel. +39 340.9657108
Fax +39 011.6647421

E-mail: elisabetta.bersezio@unito.it

The **Journal of Endocrinological Investigation** is published monthly by Editrice Kurtis s.r.l., and is cited in Current Contents/ Index Medicus, Excerpta Medica, Biosis, Chemical Abstracts, SIIC (Sociedad Iberoamericana de Información Científica).

Copyright

Manuscripts are accepted for publication with the understanding that no substantial part has been, or will be published elsewhere. By submitting a manuscript, the authors agree that the copyright is transferred to the Publisher if and when the article is accepted for publication. The copyright covers the exclusive rights to reproduce and distribute the article, including reprints, photographic reproduction and translation. No part may be duplicated or reproduced without permission of the Publisher and the Editor. Printed in Italy.

If products, dosages and methods of employment appear in this publication, the Publisher cannot grant their exactness and suggests that the reader should refer to the pertinent Literature.

Induction of vascular endothelial growth factor release by transcutaneous frequency modulated neural stimulation in diabetic polyneuropathy

M. Bevilacqua¹, L.J. Dominguez², M. Barrella¹, and M. Barbagallo²

¹Endocrinology and Diabetes Unit and LORENZ Research Center, Department of Medicine, Luigi Sacco Hospital (Vialba)-University of Milan, Milan; ²Geriatric Unit, Department of Internal Medicine and Geriatrics, University of Palermo, Palermo, Italy

ABSTRACT. *Background:* Pharmacological treatment for diabetic polyneuropathy (DP) has shown limited benefit; frequency-modulated electrical stimulation (FREMS) has shown positive results in pain control and nerve conduction velocity in DP. *Objective:* To investigate the effects of FREMS vs transcutaneous electrical nerve stimulation (TENS) on the release of vascular endothelial growth factor (VEGF) in Type 2 diabetic and in non-diabetic subjects. *Methods:* 10 non-diabetic [mean age 37±5 yr; females (F)/males (M): 6/4] and 10 Type 2 diabetic subjects (mean age 52±6 yr; F/M: 5/5) with DP underwent TENS (for 10 min) followed by 30 min interval without electrical stimulation, and then FREMS (for 10 min) over the forearm volar surface. Blood samples for VEGF measure-

ments were obtained from the contra-lateral arm every 2 min during TENS/FREMS application and every 10 min during the intervals. *Results:* We observed a significant rise in plasma VEGF during FREMS in both non-diabetic and diabetic subjects (maximal response 89.4±80.3 pg/ml and 48.5±18.3 pg/ml, respectively; $p<0.01$ vs basal) with a lower, but still significant response in diabetics. No changes in VEGF were observed during TENS application. *Conclusion:* VEGF release during FREMS may help explain the positive effects on nerve conduction velocity in DP, possibly mediated by favorable effects on *vasa nervorum* microangiopathy.

(J. Endocrinol. Invest. 30: 944-947, 2007)

©2007, Editrice Kurtis

INTRODUCTION

Peripheral neuropathy is a frequent complication of long-term diabetes that ultimately accounts for considerably disabling morbidity (1, 2). Clinical hallmarks of diabetic polyneuropathy (DP) are chronic pain and impaired nerve conduction velocity in both, motor and sensory pathways, while histopathological features include *vasa nervorum* microangiopathy, and axons atrophy/loss resulting from different tissue damage mechanisms common to all long-term diabetic complications (1). Final consequences of such damage include foot ulcerations, accounting for ~20% of hospital admissions among diabetics (3), which may lead to lower ex-

tremity amputation, three times more common in diabetic compared to non-diabetic subjects (4). Proposed pathogenetic mechanisms for DP include metabolic injury to nerve fibers (5), *vasa nervorum* damage (6), sorbitol accumulation (7), fatty acids alterations (8), protein glycosylation (9), oxidative stress with changes in NAD:NADH ratio (10), neurotrophic growth factors reduction (11), and autoimmune damage (12). This multifactorial process lead to progressive nervous fiber demyelination, functional deterioration and suppressed regeneration (1, 2). Gene transfer of plasmid DNA encoding vascular endothelial growth factor (VEGF) has been studied in experimental diabetic animals with positive results (13). Induction of VEGF synthesis and release by sub-contraction electrical stimulation in skeletal muscle cells *in vitro* and in experimental hind limb ischemia *in vivo* has been reported (14). Current evidence have demonstrated the central role that glycemic tight control and reduction of other cardiovascular risk factors may have on DP development and progression (2), but not in the modification of the disease once established. Phar-

Key-words: Diabetic neuropathy, diabetes mellitus, VEGF, electrical stimulation, TENS.

Correspondence: M. Bevilacqua, MD, Servizio di Endocrinologia e Diabetologia, Ospedale L. Sacco, Via GB Grassi 74, 20157 Milan, Italy.

E-mail: maurziobevilacqua@fastwebnet.it

Accepted August 18, 2007.

macological trials for DP have shown limited benefit, hence, some attention has been given to non-pharmacological approaches. Transcutaneous electrical nerve stimulation (TENS) showed benefit in pain relief (15) and, recently, electrical transcutaneous stimulation with modulated frequencies (FREMS) has demonstrated positive effects not only on pain control, but also on nerve conduction velocity (16). The present study aimed to investigate the effects of FREMS compared to TENS on VEGF release in diabetic and non-diabetic subjects.

SUBJECTS AND METHODS

Ten diabetic patients [mean age 52 ± 6 yr; females (F)/males (M): 5/5] with DP were recruited from the Diabetes Outpatient Clinic and 10 nondiabetic subjects (mean age 37 ± 5 yr; F/M: 6/4) were recruited from our department's employees who volunteered to participate in the study after being carefully informed about the purpose and the procedures involved. All participants gave informed consent, the study was approved by the local Ethics Committee and conducted in accordance with the Declaration of Helsinki. All subjects underwent TENS for 10 min, followed by an interval of 30 min without electrical stimulation. Then FREMS therapy was applied for 10 min. Both electrotherapies were administered with APTIVA neurostimulator (Lorenz Therapy™ System, Medolla, MO, Italy) via 2 non-woven electrodes applied over the forearm volar surface. DP was identified with electromyography (Medelec™ Synergy N-EP-EMG/EP Monitoring System). According to international criteria and standardization at our institution, two of the following criteria, bilaterally, were considered for DP diagnosis (mean of 6 speed conduction for each nerve): 1) ulnar motor conduction velocity (MCV) < 48 m/sec, sensory conduction velocity (SCV) < 50 m/sec; 2) common peroneal MCV < 45 m/sec and/or amp < 3 mV; 3) posterior tibialis MCV < 42 m/sec and/or amp < 4 mV; 4) suralis SCV < 48 m/sec and/or amp < 8 mV.

FREMS consisted of sequences of mono-phase compensated negative potential electric pulses, with sharp spike and asymmetrical shape; peak amplitude varied from 0 to 255 Volts; pulse frequency varied from 1-50 Hz; pulse duration varied from 10-40 microseconds. Blood samples for plasma VEGF measurements (enzyme-linked immunosorbent assay; R&D Systems) were obtained from the contralateral arm through intravenous indwelling catheter at different time points (indicated in Fig. 1) during 70 min. Statistical analysis was performed using GraphPad software version 4.0 (GraphPad, San Diego, CA, USA), considering statistical significance for *p* values below 0.05.

RESULTS

Plasma levels of VEGF were obtained before, during, and after both electrostimulation therapies. The mean VEGF concentration \pm SD obtained in non-diabetic subjects (upper panel) and Type 2 diabetic patients with DP (lower panel) are shown in Figure 1. FREMS treatment resulted in maximum VEGF concentrations during the 10 min applica-

tion, followed by a decrease of VEGF concentrations towards the baseline levels in both non-diabetic and diabetic subjects (Fig. 1) ($p < 0.001$ according to analysis of variance for repeated measures). The VEGF elevation observed in correspondence to FREMS administration was not observed during TENS application in both, non-diabetic and diabetic subjects. Indeed, VEGF concentrations during TENS were similar to those observed during the intervals before application of the electrostim-




Fig. 1 - Plasma vascular endothelial growth factor (VEGF) levels during (transcutaneous electrical nerve stimulation) TENS (0-10 min) and during frequency-modulated electrical stimulation (FREMS) (40-50 min) administration in non-diabetic subjects (panel A) and in Type 2 diabetic subjects with polyneuropathy (panel B). There was a significant increase of serum VEGF concentration during FREMS application in both groups, with no significant increase in VEGF concentrations during TENS application. *means $p < 0.001$.

ulation therapies. The areas under the curve for VEGF release, calculated with the trapezoidal rule, were as follows: in non-diabetics 490.1 ± 171.8 vs 246.2 ± 98.3 pg/ml during 10 min for FREMS vs TENS application, respectively ($p < 0.01$); in diabetics, 400.6 ± 124.8 vs 122.1 ± 41.0 pg/ml during 10 min for FREMS vs TENS application, respectively ($p < 0.001$). Although maximal responses were lower in diabetics (48.5 ± 18.3 pg/ml; $p < 0.01$ vs basal), compared to the maximal responses in non-diabetic subjects (89.4 ± 80.3 pg/ml; $p < 0.01$ vs basal) they were still present and significant.

DISCUSSION

Despite recent evidence that not only glycemic tight control but also the control of hypertension, overweight, high triglyceride levels, and smoking may influence the development of DP (2), few evidence-based strategies demonstrating improvement or prevention of this devastating condition are available. Controlled pharmacological trials for DP proved heretofore disappointing even with initial encouraging reports. Hence, an urgent need to develop new therapeutical approaches that improve nerve function in diabetic patients is warranted.

Non-pharmacological therapeutical attempts tested up to now include TENS, which showed some benefit in pain reduction (15), but without affecting nerve conduction velocity. We and others have used electro-magnetic neural stimulation with frequency modulation, FREMS, in patients with DP. FREMS therapy induces both, pain relief and significant increases in motor nerve conduction velocity in these patients (16).

An experimental study tested the hypothesis that DP results from *vasa nervorum* destruction and that it can be reversed by administration of angiogenic growth factors. Rats with streptozotocin-induced diabetes and rabbits with alloxan-induced diabetes in whom nerve blood flow and vessel number were markedly attenuated, paralleled to significant slowing of motor and sensory nerve conduction velocities, were compared to control non-diabetic animals. Intramuscular gene transfer of plasmid DNA encoding VEGF-1 or VEGF-2 induced revascularization and blood flow restoration in nerves of treated animals to levels of those observed in non-diabetic controls; constitutive overexpression of both transgenes resulted in restoration of large and small fiber peripheral nerve function (17). These results support the notion that DP results from *vasa nervorum* ischemia, hence, VEGF is proposed as a candidate treatment for this condition.

VEGF, the most potent endogenous angiogenic factor so far identified, is produced in cells around the vasculature and acts selectively on vascular endothelial cells as a paracrine factor stimulating angiogenesis both *in vitro* and *in vivo*. Expression of VEGF is highly regulated by hypoxia, providing a feedback mechanism to improve oxygenation via the promotion of new vessel formation (2). Kanno et al. showed the induction of VEGF protein synthesis and release by sub-contraction electrical stimulation in skeletal muscle cells in culture. This effect was confirmed *in vivo* in an experimental model of hind limb ischemia, in which increased muscle blood flow and VEGF synthesis was observed after electrical stimulation (14).

The difference in VEGF release observed in the present study with two electrotherapies may reside in the variability of electrical stimulus frequency and voltage amplitude induced by FREMS, but not by TENS. The modulation of these parameters induces composite Motor Action Potentials (MAP) in excitable tissues by the summation of sub-threshold electric stimulations, conveyed through the skin over the motor nerve. A single, low-intensity and brief duration impulse, such as the one delivered by TENS, is unable to overcome the dielectric skin barrier and excite the underlying nervous and/or muscle to elicit a recordable MAP; this effect can be achieved by FREMS through specific sequences of weak impulses with rapid increases and decreases in pulse frequency and duration, which may result in gradual recruitment of MAP in the stimulated tissues (18). MAP induction in the underlying neural tissue may have stimulated endothelium of *vasa nervorum* to release VEGF into the circulation. An improvement in endoneural blood flow, hypoxia, and microvessel disease may help to explain the positive reported effects of FREMS on DP (16). In fact, blood flow improvement with either revascularization or physical exercise is accompanied with increased nerve conduction velocity. Studies on the chronic effect of FREMS on VEGF synthesis and release are needed in order to clarify the nature of long-term FREMS actions.

REFERENCES

1. DUBY JJ, CAMPBELL RK, SETTER SM, WHITE JR, RASMUSSEN KA. Diabetic neuropathy: an intensive review. *Am J Health Syst Pharm* 2004, 61: 160-73.
2. TESFAYE S, CHATURVEDI N, EATON SEM, et al; EURODIAB Prospective Complications Study Group. Vascular risk factors and diabetic neuropathy. *N Engl J Med* 2005, 352: 341-50.

3. Struijs JN, Baan CA, Schellevis FG, Westert GP, van den Bos GA. Comorbidity in patients with diabetes mellitus: impact on medical health care utilization. *BMC Health Serv Res* 2006, 6: 84.
4. Lavery LA, Wunderlich RP, Tredwell JL. Disease management for the diabetic foot: effectiveness of a diabetic foot prevention program to reduce amputations and hospitalizations. *Diabetes Res Clin Pract* 2005, 70: 31-7.
5. Cameron NE, Cotter MA. Metabolic and vascular factors in the pathogenesis of diabetic neuropathy. *Diabetes* 1997, 46 (Suppl 2): S31-7.
6. Veves A, King GL. Can VEGF reverse diabetic neuropathy in human subjects? *J Clin Invest* 2001, 107: 1215-8.
7. Greene DA, Lattimer SA, Sima AA. Are disturbances of sorbitol, phosphoinositide, and Na⁺-K⁺-ATPase regulation involved in pathogenesis of diabetic neuropathy? *Diabetes* 1988, 37: 688-93.
8. Horrobin DF. Essential fatty acids in the management of impaired nerve function in diabetes. *Diabetes* 1997, 46 (Suppl 2): S90-3.
9. Brownlee M. Glycation products and the pathogenesis of diabetic complications. *Diabetes Care* 1992, 15: 1835-43.
10. Obrosova IG. How does glucose generate oxidative stress in peripheral nerve? *Int Rev Neurobiol* 2002, 50: 3-35.
11. Apfel SC, Arezzo JC, Brownlee M, Federoff H, Kessler JA. Nerve growth factor administration protects against experimental diabetic sensory neuropathy. *Brain Res* 1994, 634: 7-12.
12. Granberg V, Ejskjaer N, Peakman M, Sundkvist G. Auto-antibodies to autonomic nerves associated with cardiac and peripheral autonomic neuropathy. *Diabetes Care* 2005, 28: 1959-64.
13. Shweiki D, Itin A, Soffer D, Keshet E. Vascular endothelial growth factor induced by hypoxia may mediate hypoxia-initiated angiogenesis. *Nature* 1992, 359: 843-5.
14. Kanno S, Oda N, Abe M, et al. Establishment of a simple and practical procedure applicable to therapeutic angiogenesis. *Circulation* 1999, 99: 2682-7.
15. Kumar D, Marshall HJ. Diabetic peripheral neuropathy: amelioration of pain with transcutaneous electrostimulation. *Diabetes Care* 1997, 20: 1702-5.
16. Bosi E, Conti M, Vermigli C, et al. Effectiveness of a novel frequency modulated electro-magnetic neural stimulation in the treatment of painful diabetic neuropathy. *Diabetologia* 2005, 48: 817-23.
17. Schratzberger P, Walter DH, Rittig K, et al. Reversal of experimental diabetic neuropathy by VEGF gene transfer. *J Clin Invest* 2001, 107: 1083-92.
18. Barrella M, Toscano R, Goldoni M, Bevilacqua. Frequency rhythmic electrical modulation system (FREMS) on H-reflex amplitudes in healthy subjects. *Eura Medicophys* 2007, 43: 37-47.

INSTRUCTIONS TO AUTHORS

The *Journal of Endocrinological Investigation* publishes clinical and experimental studies from all areas of endocrinology, metabolism and andrology.

MANUSCRIPTS

The following types of manuscript are published in the *Journal of Endocrinological Investigation*: Original Papers, Case Reports, Comments, Review Articles, Short Reviews, Editorials, Opinions, Rapid Communications, Letters and Clinical Symposia. Original Papers should be based on original rather than confirmatory data, and should conform to the organizational plan outlined below. Case Reports should illustrate either a previously unrecognized disorder or a new aspect of a known condition. Comments are reserved to concise articles that are primarily based on essentially negative or confirmatory data or are of a relatively minor relevance. Review and Short Review articles on selected topics by authorities in the field are published. Both Reviews and Short Reviews are in general invited by the editors but suggestions by interested individuals may also be considered. Prospective authors should submit a formal and detailed proposal to the Editor, indicating the title and a brief outline of the content. Manuscripts should provide an up-to-date and authoritative review of a topic in any area of experimental and clinical endocrinology, metabolism and andrology. Invited Editorials linked to recent high-quality papers are also published. Opinions written by recognized authorities in their own field of research will be published only under direct invitation by the Editorial Board. Rapid Communications are reserved to novel findings that require fast publication, while confirmatory or negative data will not be considered for this kind of publication. Letters providing pertinent comments on published articles will be considered and the authors concerned will be given a right to reply. Letters raising problems of general interest in endocrinology, metabolism and andrology will also be considered. Publication of correspondence is at the discretion of the Editorial Board. Clinical Symposia from invited contributors are published occasionally. Special Issues devoted to a specific topic will be published once or twice a year.

SUBMISSION PROCEDURES

Submission of manuscripts implies that no substantial part has been, or will be published elsewhere and a Copyright Agreement should be included with all submitted manuscripts. Manuscripts are published in the order of final acceptance after review, revision and return of approved galley proofs.

The *Journal of Endocrinological Investigation* publishes manuscripts received only by online submission. Authors should submit their manuscripts through <http://jei.manuscriptcentral.com>.

Authors should prepare their manuscript as double-spaced MS Word (DOC) files to be uploaded and automatically converted into PDF format. Word 2000 (PC), Word 1998 (MAC) and previous versions are accepted.

The system will accept standard image formats and converts them to JPEG file format. All figures may be uploaded separately or embedded within the source manuscript file.

Further information will be available online at the web site.

MANUSCRIPT ORGANIZATION AND STYLE

Manuscripts must be written in concise and correct English. They should be carefully checked for typing, spelling and grammar before they are submitted. Manuscripts should be organized as

follows: Title Page; Abstract; Text; Acknowledgments; References; Figure Legends and Tables.

Title Page

This should include in order:

Title of paper. The title should be concise and clear, and should not contain abbreviations.

Authors (Initials of the name and surnames). The name of the last Author should be preceded by "and". The Affiliations (in English) should be attributed to the Authors using apex numbers. E.g.:

S. Smith¹, M. Rossi² and R. Green¹

¹Division of ..., Department of ..., University of ..., City, Country;

²Division of ..., Department of ..., University of ..., City, Country.

abbreviated version of the Title not exceeding 40 typewritten characters.

a list of 5 key-words for indexing purposes.

Name, Degree (e.g. S. Smith, M.D., Ph.D.) and full Address - Telephone - Fax - E-mail of the Corresponding Author to whom the proofs and requests for reprints should be sent.

Acknowledgements:

List of the abbreviations (see below)

Abstract

The text must be preceded by a structured abstract of no more than 250 words to include the following headings: Background, Aim, Material/Subjects and Methods, Results, Conclusions. The abstract should be self-explanatory, without reference to the text. Abbreviations may be included, provided that they are defined in the abstract as well as in the main text.

Main text

Original articles should conform to the following organizational plan: Introduction - Materials and Methods - Results - Discussion. The Introduction should include a brief review of the literature data strictly related to the object of the paper, and a short statement on the aims of the work presented.

Materials and Methods should be described with sufficient details to allow others to duplicate the study. Previously reported procedures may be referred to by citation, but newly adopted modifications should be specified in detail.

Results must be clearly and concisely described with the help of appropriate illustrative material. Results should be expressed in International System Units or the conversion factor to the International System Units should be clearly indicated. Time of day should be expressed using the 24-hour clock (e.g., 08:00 h, 15:00 h).

The Discussion should be limited to the reported findings and their implications.

Footnotes to the text should be typed and numbered consecutively at the foot of the appropriate page, using a line to separate from the text.

References

The number of References should be adequate, but not excessive; to this purpose, the use of appropriate recent reviews is recommended. Each reference should be cited in numerical order (in parentheses) in the text and listed in the same numerical

order at the end of the paper. Articles in press (i.e. accepted for publication) may be included in the bibliography; the name of the journal in which they will appear, and if possible volume and year, should be indicated. References to unpublished material may be cited in parentheses in the text, but not in the bibliography.

Abstracts may be cited only when they contain substantial data not published elsewhere, their nature should be stated in the bibliography by the addition of the term "(abstract)" at the end of the reference. Authors should check references against original sources for accuracy.

References to the literature should be listed indicating all the Authors when six or less than six, or only the first three Authors followed by "et al." when more than six in total.

The titles of journals should be abbreviated according to the style used in the Index Medicus.

Neither bold nor italics characters should be present in the Reference list.

Sample References

Example of reference to article in Journals with more than 6 Authors:

Levin SR, Hofeldt FD, Schneider V, et al. Cryohypophysectomy for acromegaly. Factors associated with altered endocrine function and carbohydrate metabolism. *Am J Med* 1974, 57: 526-35.

Example of reference to article in Journals with 6 Authors or less:
Charro AL, Hofeldt FD, Becker N, Levin SR, Forsham PH. Adrenocortical function in acromegaly. *Am J Med Sci* 1973, 266: 211-8.

Example of reference to abstracts (List all Authors if 6 or less; only the first 3 Authors followed by "et al." when more than 6 in total):

MacLaughlin DT, Cigarros F, Donahoe PK. Mechanism of action of Mullerian inhibiting substance. Program of the 70th Annual Meeting of The Endocrine Society, New Orleans, LA, 1988, p. 19 (abstract).

Example of reference to books (List all Authors if 6 or less; only the first 3 Authors followed by et al. when more than 6 in total)
Bonnevillie F, Cattin F, Dietemann JL. Computed tomography of the pituitary gland. Heidelberg: Springer-Verlag 1986, 15-164.

Burrow GN. The thyroid: nodules and neoplasia. In: Felig P, Baxter JD, Broadus AE, Frohman LA eds. *Endocrinology and metabolism*. 2nd ed. New York: McGraw-Hill. 1989, 473-507.

Illustrations

Figures should be numbered consecutively with arabic numbers. Each figure must have a concise legend. Color illustrations will be published when approved by the Editorial Board at the Authors' expense.

Tables

Tables should be numbered consecutively using arabic numbers. Each table must have a concise heading and should be comprehensible without reference to the text. Tabular data should in general not be duplicated in the text or in the figures.

RAPID COMMUNICATIONS

Manuscripts must strictly adhere to the JEI Instructions to Authors regarding the general organization of the paper. Manuscripts

should not exceed 15,900 typewritten characters from Title to References included. One Table and one Figure (or 2 Tables or 2 Figures) sized 78 mm wide and 61 mm high or 163 mm wide and 86 mm high are allowed. For each figure or table sized 78x61 mm, 1000 typewritten characters should be subtracted from the maximum allowed number of typewritten characters. For each figure or table sized 163x86 mm, 2500 typewritten characters should be subtracted from the maximum allowed number of typewritten characters.

OPINIONS

Manuscripts should have no more than 2 Authors and should not exceed 15,900 typewritten characters from Title to References included. No more than 2 Authors and 15 references should be included. One Table and one Figure (or 2 Tables or 2 Figures) sized 78 mm wide and 61 mm high or 163 mm wide and 86 mm high are allowed. For each figure or table sized 78x61 mm, 1000 typewritten characters should be subtracted from the maximum allowed number of typewritten characters. For each figure or table sized 163x86 mm, 2500 typewritten characters should be subtracted from the maximum allowed number of typewritten characters.

EDITORIALS

Manuscripts should have no more than 2 Authors and should not exceed 10,600 typewritten characters from Title to References included. No more than 5 references, including the paper to which the Editorial is linked, are allowed. Editorials do not contain illustrations.

SHORT REVIEWS

Short review articles are welcome on late breaking discoveries on which a substantial literature has not yet been accumulated. A Short Review should provide a forum for sharply focused topics of recent experimental or clinical investigations in medicine primarily to make these results accessible to both clinical and basic researchers. Short Reviews may also reflect the state-of-the-art of the field being reviewed and often include a short history, although there is no need to include an exhaustive literature survey. Manuscripts should not exceed 31,800 typewritten characters from Title to References included. No more than 50 references should be included.

PUBLICATION CHARGES

Authors will be charged € 54,00 for each printed page including tables, graphs and illustrations. Authors will also be requested to contribute to the cost of color illustrations.

GALLEY PROOFS

Galley proofs will be sent to the Corresponding Author, unless otherwise indicated. Proofs should be returned within 48 hours from receipt with clear indication of any changes.

REPRINTS

Reprints may be purchased at the prices indicated on the reprint order form accompanying the galley proofs.

REVIEWING

All submitted manuscripts are examined by the Editorial Board and sent to at least two expert Referees. Authors are usually notified within 4 to 6 weeks about the acceptability of their manuscript.